

Rosiński Measures for Tempered Stable and Related Ornstein-Uhlenbeck Processes*

György TERDIK[†] and Wojbor A. WOYCZYŃSKI[‡]

September 5, 2006

Abstract

Several concrete parametric classes of tempered stable distributions are discussed in terms of explicit calculations of their *Rosiński measures*. The hope that they will provide a family of concrete models useful in applied areas and for which the fitting can be done by parametric methods. Related Ornstein-Uhlenbeck processes are analyzed. The emphasis throughout the paper is on obtaining exact analytic formulas.

Dedicated to the memory of Kazimierz Urbanik

Keywords: stable distributions, Ornstein-Uhlenbeck and Lévy processes, Truncated Lévy Flights.

Subject Classification: 62M15, 60G60

1 Introduction

Ever since the pioneering applied work by Mandelbrot [12], and Montroll and Scher [15], the Lévy stable processes enjoyed a great popularity as a flexible modeling tool in natural and economic sciences. However, their elegant scaling properties which made them analytically pleasing were also a problem when people tried to fit them to real data; the scaling could not be maintained at all scales, and the thick tail behavior was often impossible to verify rigorously. The remedy proposed first by Mantegna and Stanley [13], and others, was to keep the local behavior of the distributions Lévy-stable like but to truncate the tails. Different schemes for the truncation and their parametrizations were proposed and discussed in the physical literature (see, e.g., [27] for a partial history of these efforts). All of those schemes suffered, however, from the lack of invariance under linear transformations for the distributions classes introduced, a property that Lévy stable distributions enjoyed. Such was the state of affairs until publication of Rosinski's paper [18] where a formal and elegant definition of the class of *tempered stable distributions and processes* was proposed. The latter did have invariance under the linear transformations is one of their most important structural properties.

*This work has been partly carried out while Gy. T. was visiting at the Department of Statistics and Center for Stochastic and Chaotic Processes in Science and Technology, Case Western Reserve University, Cleveland, OH 44106. It was partially supported by the Hungarian NSF OTKA No. T047067 (Gy.T), and by the U.S. National Science Foundation (W.A.W.).

[†]Department of Inf. Techn., University of Debrecen, 4010 Debrecen, Hungary, e-mail: terdik@delfin.unideb.hu .

[‡]Department of Statistics and Center for Stochastic and Chaotic Processes in Science and Technology, Case Western Reserve University, Cleveland, OH 44106, e-mail: waw@case.edu

The class of tempered stable distributions has in infinite-dimensional parametrization by a family of measures, which makes their fitting to real data a difficult task. For this reason, we embark in this paper on a fairly pedestrian project of developing a series of parametric models that fit into the general framework of tempered distributions but for which parametric statistical estimation procedures can be realistically developed. The emphasis is on obtaining explicit analytic formulas and on explicit calculations. Once those distributional models are developed it is natural to study the corresponding Lévy processes and, more physically meaningful, corresponding Ornstein-Uhlenbeck processes.

The paper begins, in Section 2, with brief preliminaries on infinitely divisible distributions in \mathbb{R}^d , and the subclass of self-decomposable distributions which were developed, among others, by Urbanik, see, [28]–[30]. Here, the role of cumulants and cumulant functions, which will be paramount throughout the paper, is first explained.

Section 3 begins with the basic definitions of tempered stable distributions and introduces the concept of the Rosiński measure (R-measure). A subsection on fundamental properties of tempered stable distributions follows. The section ends with a discussion of the relation of this class, in one- and two-dimensional settings, to the previously studied smoothly truncated Lévy distributions. It is Section 4 where we introduce and discuss several parametric tempering schemes using a variety of special functions: gamma, inverse Gaussian, fractional exponential, 1/3-stable and Bessel.

Finally, in Section 5 we turn to the multivariate tempered stable Ornstein-Uhlenbeck processes, their background driving Lévy processes (BLDP), and their stationary versions. An appendix on the multiple cumulant technique, not commonly seen in the literature, concludes the paper.

2 Preliminaries

Let us begin with a random variable X having an infinitely divisible distribution P on \mathbb{R}^d , with a generating triple (Σ, M, b) , and the characteristic function given by

$$\varphi(y) = \exp\left(-\frac{1}{2}\langle y, \Sigma y \rangle + i\langle b, y \rangle + \int_{\mathbb{R}^d} \left(e^{i\langle y, x \rangle} - 1 - i\langle y, x \rangle \mathbf{1}_D(x)\right) M(dx)\right),$$

where $D = \{x : \|x\| \leq 1\}$, and M is a Lévy measure on \mathbb{R}^d , satisfying

$$M(\{0\}) = 0, \quad \text{and} \quad \int_{\mathbb{R}^d} (\|x\|^2 \wedge 1) M(dx) < \infty.$$

Furthermore, throughout this paper, we are going to assume that

$$\int_{\|x\|>1} \|x\|^m M(dx) < \infty, \tag{1}$$

for some $m \geq 1$. This condition is equivalent (see, Sato [20], Theorem 25.3) to the condition

$$E \|X\|^m < \infty. \tag{2}$$

If, additionally,

$$\int_{\|x\|<1} \|x\|^m M(dx) < \infty, \tag{3}$$

then we can rewrite the characteristic function $\varphi(y)$ in the centered form (cf., [20], p. 39)

$$\begin{aligned}\varphi(y) &= \exp \left(iy^\top c_1 + i^2 \frac{1}{2!} y^\top \otimes^2 c_2 + \cdots + i^k \frac{1}{k!} y^\top \otimes^m c_m + \int_{\mathbb{R}^d} \left(e^{i\langle y, x \rangle} - \sum_{j=0}^m \frac{i^j}{j!} y^\top \otimes^j x^{\otimes j} \right) M(dx) \right) \\ &= \exp \left(\sum_{j=1}^m \frac{i^j}{j!} y^\top \otimes^j c_j + \int_{\mathbb{R}^d} \left(e^{i\langle y, x \rangle} - \sum_{j=0}^m \frac{i^j}{j!} y^\top \otimes^j x^{\otimes j} \right) M(dx) \right),\end{aligned}$$

where c_j is the j^{th} order cumulant of the random variable X , that is, the "moment" of order j of the Lévy measure M , and \otimes , and \top , stand, respectively, for the Kronecker product, and the transpose. Recall that if A is an $m \times n$ -matrix, and B is a $p \times q$ matrix, then the Kronecker product $A \otimes B$ is the $mp \times nq$ block matrix

$$A \otimes B = \begin{bmatrix} a_{11}B & \cdots & a_{1n}B \\ \vdots & \cdots & \vdots \\ a_{m1}B & \cdots & a_{mn}B \end{bmatrix}.$$

If $m \geq 1$, then c_1 is the center of the measure P ; if $m \geq 2$ then c_2 is the vector of the variances of X . Note that c_1 , and c_2 , are not uniquely determined by the Lévy measure M unless the distribution P has no Gaussian component ($\Sigma = 0$) and is centered ($c_1 = 0$).

The distribution P is said to be *self-decomposable* if, for every $\gamma > 1$, there exists a probability measure P_γ on \mathbb{R}^d such that

$$\varphi(y) = \varphi(\gamma^{-1}y) \varphi_\gamma(y),$$

where φ_γ is the characteristic function of P_γ . A self-decomposable P is necessarily infinitely divisible and, for any $\gamma > 1$, the measure P_γ is a uniquely determined infinitely divisible measure. The infinitely divisible distribution P is a self-decomposable if, and only if, its Lévy measure M is of the form

$$M(B) = \int_{\mathbb{S}^{d-1}} \int_0^\infty 1_B(ru) k_u(r) \frac{dr}{r} \sigma(du), \quad (4)$$

where \mathbb{S}^{d-1} is the unit sphere in \mathbb{R}^d , $\sigma(du)$ is a finite measure on \mathbb{S}^{d-1} , and $k_u(r)$ is decreasing in r , and measurable in u . In particular, for $d = 1$, the characteristic function of a self-decomposable measure is of the form

$$\varphi(y) = \exp \left(-\frac{1}{2} \Sigma y^2 + iby + \int_{-\infty}^\infty (e^{iyx} - 1 - iyx 1_{[-1,1]}(x)) k(x) \frac{dx}{|x|} \right),$$

where $\Sigma \geq 0$, $b \in \mathbb{R}$, and $k(x)$ is a nonnegative function, increasing on $(-\infty, 0)$, decreasing on $(0, \infty)$, and such that

$$\int_{-\infty}^\infty (|x|^2 \wedge 1) k(x) \frac{dx}{|x|} < \infty,$$

see [28], [30], [29], [20] p. 109..

3 Tempered Stable Distributions

3.1 Basic Definitions.

A Gaussian-free ($\Sigma = 0$), self-decomposable probability distribution is said to be *tempered α -stable* if, in the polar representation (4) of its Lévy measure M , the function

$$k_u(r) = k(r|u)r^{-\alpha},$$

where $\alpha \in (0, 2)$, and $k(\cdot|u)$ is a completely monotone function, with $k(0+|u) = 1$, and $k(\infty|u) = 0$. In other words, the Lévy measure of the tempered α -stable distribution is of the form

$$M(B) = \int_{\mathbb{S}^{d-1}} \int_0^\infty 1_B(ru) k(r|u) \frac{dr}{r^{\alpha+1}} \sigma(du), \quad (5)$$

where $\sigma(du)$ is a finite measure on the unit sphere \mathbb{S}^{d-1} . The parameter α is called the *index*, and the function k – the *tempering function*.

Obviously, the case of constant $k(r|u)$ corresponds to the classical α -stable distribution. The above concept of the tempered stable distribution is due to Rosiński [18], who made essential use of the fact that the tempering function k can be represented as the Laplace transform

$$k(r|u) = \int_0^\infty e^{-rs} Q(ds|u),$$

where $Q(ds|u)$ is a u -measurable family of probability measures on \mathbb{R}_+ . The σ -weighted superposition of the measures $Q(ds|u)$ defines a finite measure

$$Q(B) = \int_{\mathbb{S}^{d-1}} \int_0^\infty 1_B(ru) Q(dr|u) \sigma(du), \quad (6)$$

on \mathbb{R}^d . The formula

$$R(B) = \int_{\mathbb{R}_0^d} 1_B\left(\frac{x}{\|x\|^2}\right) \|x\|^\alpha Q(dx), \quad \mathbb{R}_0^d = \mathbb{R}^d \setminus \{0\}, \quad (7)$$

defines a measure, equivalent with the measure Q which is, in turn, can be expressed in terms of the measure R as follows:

$$Q(B) = \int_{\mathbb{R}_0^d} 1_B\left(\frac{x}{\|x\|^2}\right) \|x\|^\alpha R(dx).$$

Measures Q and R have been demonstrated (see, Rosiński [18], [19]) to be of importance for different applications, with Q being particularly useful for the simulation of the tempered stable random variables. We shall refer to R as the *Rosiński measure*, or *R-measure*, for short. An R-measure R together with a constant b , uniquely determine, via (4), a tempered α -stable measure. The class of all tempered α -stable measures (or, loosely speaking, related random variables) will be denoted by $TS_\alpha(R, b)$.

3.2 Properties of Tempered Stable Distributions.

We begin this subsection by listing the fundamental properties of $TS_\alpha(R, b)$.

(i) *The cumulant function representation.* The cumulant function (that is, the logarithm of the characteristic function) κ_X of a random variable X with a tempered stable distribution in $TS_\alpha(R, b)$ is uniquely given (see [18] for details) by the formula

$$\kappa_X(y) = \int_{\mathbb{R}_0^d} \psi_\alpha(\langle y, x \rangle) R(dx) + i \langle y, b \rangle, \quad (8)$$

where

$$\psi_\alpha(r) = \begin{cases} \Gamma(-\alpha) [(1-ir)^\alpha - 1], & 0 < \alpha < 1, \\ (1-ir) \log(1-ir) + ir, & \alpha = 1, \\ \Gamma(-\alpha) [(1-ir)^\alpha - 1 + i\alpha r], & 1 < \alpha < 2. \end{cases} \quad (9)$$

(ii) *Moments.* A random variable $X \in TS_\alpha(R, b)$ has always finite moments of order $m < \alpha$. If $m \geq \alpha$, then the moment assumption (1) can be expressed in terms of the R -measure as follows:

$$E \|X\|^m < \infty,$$

if and only if

$$\int_{\|x\|>1} \|x\|^m R(dx) < \infty. \quad (10)$$

If the support of R is a bounded set then some exponential moments are also finite.

(iii) *Invariance of $TS_\alpha(R, b)$ under independent summation.* If random variables X , and Y , are independent, and such that $X \in TS_\alpha(R_1, b_1)$ and $Y \in TS_\alpha(R_2, b_2)$, then $X + Y \in TS_\alpha(R_1 + R_2, b_1 + b_2)$.

(iv) *Invariance of $TS_\alpha(R, b)$ under linear transformations.* If $X \in TS_\alpha(R, b)$, and V is an $m \times d$ matrix then $VX \in TS_\alpha(R_V, b_V)$, where $R_V(B) = R \circ V^{-1}(B) = R(x | Vx \in B)$ and $b_V = Vb$.

The above invariance property can be extended to linear functionals of the tapered stable Lévy processes as follows: Let $Z_t, t \geq 0$, be a Lévy process with $Z_1 \in TS_\alpha(R_Z, 0)$, and

$$X = \int_0^\tau g(s) dZ_s,$$

with a fixed $\tau > 0$, and is a continuous (matrix-valued) function $g(s)$. Then, the cumulant function of X ,

$$\kappa_X(y) = \int_{\mathbb{R}_0^d} \psi_\alpha(\langle y, x \rangle) R_X(dx),$$

where

$$R_X(B) = \int_{\mathbb{R}_0^d} \int_0^\tau 1_B(g(s)x) ds R_Z(dx). \quad (11)$$

Thus (see , e.g., [11], [9]) the characteristic function of the linear functional X can be written in the form

$$\varphi_X(y) = \exp \left[\int_0^\tau \kappa_{Z_1}(g(s)^\top y) ds \right], \quad (12)$$

so that the cumulant function κ_X of X can be rewritten as follows:

$$\begin{aligned} \kappa_X(y) &= \int_0^\tau \kappa_{Z_1}(g(s)^\top y) ds \\ &= \int_0^\tau \int_{\mathbb{R}_0^d} \psi_\alpha(\langle g(s)^\top y, x \rangle) R_Z(dx) ds \\ &= \int_{\mathbb{R}_0^d} \int_0^\tau \psi_\alpha(\langle g(s)^\top y, x \rangle) ds R_Z(dx). \end{aligned}$$

(v) *Gamma-like limit, for $\alpha \rightarrow 0$.* For each $r \neq 0$, and $\alpha \rightarrow 0$, the limit of the cumulant function $\psi_\alpha(r)$ in (9) is easily calculated:

$$\lim_{\alpha \rightarrow 0} \psi_\alpha(r) = -\log(1 - ir).$$

So, if a family R_α of R -measures converges weakly to an R -measure R_X , as $\alpha \rightarrow 0$, then the function

$$\kappa_X(y) = - \int_{\mathbb{R}_0^d} \log(1 - i \langle y, x \rangle) R_X(dx)$$

is a cumulant function of a random variable X . If $d = 1$, and we choose a fixed $\lambda > 0$, and the family of R - measures $R_\alpha = \lambda^\alpha a \delta_{1/\lambda}$, where δ is the Dirac measure, then the corresponding limiting distribution (of X) is the Gamma distribution. Thus the definition of the class TS_α can be extended meaningfully to $\alpha = 0$.

(vi) *Simulations.* Rosiński [18] gives a series representation for random variables with distributions in $TS_\alpha(R, b)$ based on i.i.d. sequences of uniform, exponential and Q -distributed random variables. This series representation permits a convenient simulation of tempered stable Lévy processes and related Ornstein-Uhlenbeck processes.

(vii) *Cumulants.* The cumulants order greater than one of a tempered stable distribution can be calculated purely in terms of its R - measure R ; those of order one depend on the drift b as well.

Lemma 1 *Suppose that the R - measure R satisfies the moment condition*

$$\int_{\mathbb{R}_0^d} \|x\|^m R(dx) < \infty, \quad (13)$$

for an $m \geq 1$ in case of $0 < \alpha < 1$, and for an $m \geq 2$, when $1 \leq \alpha < 2$. Then the m^{th} order cumulant of the tempered stable random variable $X \sim TS_\alpha(R, b)$ is given by

$$c_m = \Gamma(m - \alpha) \mu_{R, \otimes m}, \quad (14)$$

where

$$\mu_{R, \otimes m} = \int_{\mathbb{R}_0^d} x^{\otimes m} R(dx).$$

Proof. First, note that the function ψ_α has the following series expansion

$$\begin{aligned} \psi_\alpha(r) &= \int_0^\infty (e^{irx} - 1 - irx) x^{-\alpha-1} e^{-x} dx \\ &= \int_0^\infty \left(e^{irx} - \sum_{j=0}^m \frac{i^j}{j!} r^j x^j \right) x^{-\alpha-1} e^{-x} dx + \sum_{j=2}^m \frac{i^j}{j!} r^j \Gamma(j - \alpha), \end{aligned}$$

which, for $\alpha = 1$, is understood as the limit for $\alpha \searrow 1$. The cumulant function κ_X has the form

$$\begin{aligned} \kappa_X(y) &= \Gamma(-\alpha) \int_{\mathbb{R}_0^d} \left[(1 - i \langle y, x \rangle)^\alpha - \sum_{j=0}^m \frac{i^j}{j!} \frac{\Gamma(j - \alpha)}{\Gamma(-\alpha)} \langle y, x \rangle^j \right] R(dx) \\ &\quad + i \langle y, b \rangle + \sum_{j=2}^m \frac{i^j}{j!} \mu_{R, j}(y) \Gamma(j - \alpha), \end{aligned}$$

where $\mu_{R, j}$ is the j^{th} moment of R , i.e.

$$\mu_{R, j}(y) = \int_{\mathbb{R}_0^d} \langle y, x \rangle^j R(dx) = y^{\top \otimes j} \int_{\mathbb{R}_0^d} x^{\otimes j} R(dx) = y^{\top \otimes j} \mu_{R, \otimes j},$$

Hence, the j^{th} order cumulant c_j of the random variable X is given by the formula

$$c_j = \Gamma(j - \alpha) \mu_{R, \otimes j},$$

see the Appendix, and also [26], for the definition of the multiple moments $\mu_{R, \otimes j}$. ■

The case of $m = 1$ is special. It is possible to find a random variable $X \sim TS_\alpha(R, b)$ with a finite first moment, but such that $\mu_{R, 1} = \infty$ (see Sections 4.4, 4.5). This fact explains why the condition (13) is stronger than the condition (10).

3.3 1-D Smoothly Truncated Lévy Distributions

One of the early examples of TS_α processes were the 1-D Truncated Lévy Flights introduced by Mantegna and Stanley [13] as a model for random phenomena which exhibit at small scales properties similar to those of self-similar Lévy processes, but have distributions which at large scales have cutoffs and thus have finite moments of any order. Koponen [10], building on Mantegna and Stanley's ideas defined the Smoothly Truncated Lévy Flights (STLF) which stressed the advantage of a nice analytic form. Independently, the same family of distributions was described earlier by Hougaard [8] in the context of a biological application. In this section, we discuss these special examples in the context of general tempered stable distributions.

The 1-D unit 'sphere' is the two-point set $\mathbb{S}^0 = \{\pm 1\}$, and Koponen's *Smoothly Truncated Lévy Distribution* ($STLD_\alpha(a, p_1, \lambda)$) is defined as a tempered α -stable distribution with the tempering function (see Subsection 3.1)

$$k(r|\pm 1) = k(r) = \exp(-\lambda r), \quad \lambda > 0, \quad r > 0,$$

and the measure

$$\sigma(\{-1\}) = ap_1, \quad \sigma(\{1\}) = ap_2,$$

where $a, p_1, p_2 > 0, p_1 + p_2 = 1$. In this case, the polar representation (5) takes the form

$$k(r|u) \frac{dr}{r^{\alpha+1}} \sigma(du) = \begin{cases} ap_1 \exp(\lambda r) \frac{dr}{|r|^{\alpha+1}}, & r < 0, \\ ap_2 \exp(-\lambda r) \frac{dr}{r^{\alpha+1}}, & r > 0. \end{cases}$$

The measures Q and R , (see (6) and (7), respectively) are given by the formulas

$$Q = ap_1 \delta_{-\lambda} + ap_2 \delta_\lambda, \quad \text{and} \quad R = \lambda^\alpha a (p_1 \delta_{-1/\lambda} + p_2 \delta_{1/\lambda}),$$

respectively, where δ_λ denotes the Dirac measure at λ . In other words,

$$Q(B) = ap_1 1_B(-\lambda) + ap_2 1_B(\lambda),$$

and

$$R(B) = \lambda^\alpha a [p_1 1_B(-1/\lambda) + p_2 1_B(1/\lambda)].$$

The cumulant function κ_X of X , see (8), is then given by

$$\kappa_X(u) = a\lambda^\alpha [p_1 \psi_\alpha(-u/\lambda) + p_2 \psi_\alpha(u/\lambda)] + iub, \quad (15)$$

with the function ψ_α defined in (9). Thus, for $m \geq 2$, the cumulants themselves are

$$\begin{aligned} \text{Cum}_m(X) &= a\lambda^\alpha \Gamma(m - \alpha) (p_1 (-\lambda)^{-m} + p_2 \lambda^{-m}) \\ &= a\lambda^{\alpha-m} \Gamma(m - \alpha) (p_1 (-1)^m + p_2). \end{aligned}$$

For a fixed $\lambda > 0$, as $\alpha \rightarrow 0$, the distribution $STLD_\alpha$ tends to the Gamma distribution $\Gamma(a, \lambda)$. Indeed, for $0 < \alpha < 1$, the Laplace transform ϕ_λ of $STLD_\alpha(a, 0, \lambda)$ is

$$\phi_\lambda(u) = \exp(a\lambda^\alpha \Gamma(-\alpha) [(1 + u/\lambda)^\alpha - 1]),$$

and

$$\lim_{\alpha \rightarrow 0} \exp\left(-a\Gamma(1 - \alpha) \frac{(\lambda + u)^\alpha - \lambda^\alpha}{\alpha}\right) = \exp(-a \log(1 + u/\lambda)) = (1 + u/\lambda)^{-a},$$

by the L'Hospital rule.

It is an interesting observation that, for $0 < \alpha < 1$, the smooth truncation of the stable cumulant function results also in the smooth truncation of the stable probability density itself. More precisely, if $f(x)$ is the density function of a one-sided α -stable distribution, then $g_\lambda(x) = f(x) e^{-\lambda|x|}$ is the density function of some Smoothly Truncated Lévy Distribution and, vice versa, if $g_\lambda(x)$ is the density function of a Smoothly Truncated Lévy Distribution, then $f(x) = g_\lambda(x) e^{\lambda|x|}$ is an α -stable density function. Moreover, there is a one-to-one correspondence between the α -stable distributions and the λ -equivalence classes of Smoothly Truncated Lévy Distributions. The connection is given by a linear transformation of the variable of the cumulant functions.

For example, the density function for the one-sided $1/2$ -stable distribution is the *Inverse Gaussian* p.d.f. (see, [24])

$$f(x) = c \frac{e^{-c^2/(2x)}}{x^{3/2} \sqrt{2\pi}}, \quad x > 0,$$

with the cumulant function, (see Sato [20], p.13)

$$\kappa(y) = -c|y|^{1/2} (1 - i \operatorname{sign}(y)).$$

The corresponding density of the Smoothly Truncated Lévy Distribution, ($\alpha = 1/2$) is

$$g_\lambda(x) = c \frac{e^{-c^2/(2x) - \lambda x + c\sqrt{2\lambda}}}{x^{3/2} \sqrt{2\pi}}, \quad x > 0,$$

and it has the cumulant function

$$\kappa_\lambda(y) = -c\sqrt{2} \left[\sqrt{\lambda - iy} - \sqrt{\lambda} \right],$$

which, after substitution $c = 2a\sqrt{\pi}$, gives the representation (15) with $\alpha = 1/2$.

Fractional and multiscaling properties of $STLD_\alpha$ have been described in [27]. In particular, we have shown that, for a one-sided distribution in $STLD_\alpha(a, 0, \lambda)$ distribution, moments of any positive order ϱ (including fractional) have the asymptotics

$$\log \mathbf{E}(|X|^\varrho) \sim \begin{cases} \min(\varrho/\alpha, 1) \log a + c_1, & \text{as } a \rightarrow 0; \\ \varrho \log a + c_2, & \text{as } a \rightarrow \infty. \end{cases}$$

For the symmetric distribution in $STLD_\alpha(a, 1/2, \lambda)$,

$$\log \mathbf{E}(|X|^\varrho) \sim \begin{cases} \min(\varrho/\alpha, 1) \log a + C_1, & \text{for } a \rightarrow 0, \\ (\varrho/2) \log a + C_2, & \text{for } a \rightarrow \infty, \end{cases}.$$

The above-quoted asymptotic results establish the multiscaling properties of the $STLD$ s and the related Smoothly truncated Lévy Flights.

3.4 2-D Smoothly Truncated Lévy Distributions

Discrete tempering. Let $X = (X_1, X_2)$ be a tempered stable random variable in \mathbb{R}^2 . Here, the unit 'sphere' \mathbb{S} consists of the vectors $u_\vartheta = (u_1, u_2) = (\cos \vartheta, \sin \vartheta)$ and, for fixed elements $u_{\vartheta_1}, u_{\vartheta_2}, \dots, u_{\vartheta_l} \in \mathbb{S}$, and constants $\lambda_1, \lambda_2, \dots, \lambda_l > 0$, we define a *Smoothly Truncated Lévy Distribution* on \mathbb{R}^2 , with index α , via the tempering function

$$k(r|u) = \exp(-\lambda_j r), \quad \text{for } u = u_{\vartheta_j}, j = 1, 2, \dots, l,$$

and zero otherwise, and a discrete measure

$$\sigma(\{e_{\vartheta_j}\}) = ap_j,$$

where $a, p_j > 0$, and $\sum p_j = 1$. The measure Q , see (6), is then

$$Q = a \sum_{j=1}^l p_j \delta_{\lambda_j e_{\vartheta_j}},$$

and R -measure, see (7),

$$R = \sum_{j=1}^l \lambda_j^\alpha p_j \delta_{1/\lambda_j e_{\vartheta_j}}.$$

The cumulant function κ_X , see (8), has now the representation

$$\kappa_X(y_1, y_2) = a \sum_{j=1}^l \lambda_j^\alpha p_j \psi_\alpha([y_1 \cos \vartheta_j + y_2 \sin \vartheta_j] / \lambda_j) + i \langle y, b \rangle.$$

with the cumulants

$$\text{Cum}_{m,n}(X_1, X_2) = a \Gamma(m+n-\alpha) \sum_{j=1}^l \lambda_j^{\alpha-m-n} p_j (\cos \vartheta_j)^m (\sin \vartheta_j)^n,$$

Remark 1. A 1-D projection of X , given by the formula $Y = v_1 X_1 + v_2 X_2$, can be viewed as a Generalized Smoothly Truncated Lévy Distribution on \mathbb{R} . Its cumulant function

$$\kappa_Y(y) = a \sum_{j=1}^l \lambda_j^\alpha p_j \psi_\alpha(yv \cos(\vartheta'_j) / \lambda_j) + i \langle y, v \rangle.$$

where $\vartheta'_j = \vartheta_j - \arctan(v_2/v_1)$, and $v = \sqrt{v_1^2 + v_2^2}$. A particular case may be obtained by matching constants v_j to $\cos(\vartheta_j)$, and constants λ_j , to obtain a cumulant function of the form

$$\kappa_Y(y) = a \sum_{j=1}^l \lambda_j^\alpha p_j \psi_\alpha(yv_j) + i \langle y, v \rangle,$$

with the cumulants

$$\text{Cum}_m(X) = a \Gamma(m-\alpha) \sum_{j=1}^l \lambda_j^{\alpha-m} p_j v_j^m.$$

Continuous tempering. Let $X = (X_1, X_2)$ be a tempered stable random variable on \mathbb{R}^2 , and $\sigma(d\vartheta) = g(\vartheta) d\vartheta$ be a finite measure on $[0, 2\pi)$. Define the tempering function k by the formula

$$k(r|\vartheta) = \exp(-\lambda(\vartheta)r)$$

where $\lambda(\vartheta) > 0$, and $r > 0$. Now, the polar representation (5) of k has the form

$$k(r|\vartheta) \frac{dr}{r^{\alpha+1}} \sigma(d\vartheta) = \exp(-\lambda(\vartheta)r) \frac{dr}{r^{\alpha+1}} g(\vartheta) d\vartheta,$$

hence

$$Q(ds|\vartheta) = \delta_{\lambda(\vartheta)}.$$

Denoting $\theta_y = \arctan(y)$, the measures Q and R are:

$$\begin{aligned} Q(B) &= \int_{\mathbb{R}_0^2} 1_B(\lambda(\theta_{x_2/x_1})x/\|x\|)g(\theta_{x_2/x_1})dx, \\ R(B) &= \int_{\mathbb{R}_0^2} 1_B(\lambda^{-1}(\theta_{x_2/x_1})x/\|x\|)\lambda(\theta_{x_2/x_1})^\alpha g(\theta_{x_2/x_1})dx. \end{aligned}$$

The cumulant function now is

$$\kappa_X(y_1, y_2) = \int_0^{2\pi} \lambda(\vartheta)^\alpha \psi_\alpha([y_1 \cos \vartheta + y_2 \sin \vartheta]/\lambda(\vartheta))g(\vartheta)d\vartheta$$

with cumulants

$$\text{Cum}_{m,n}(X_1, X_2) = \Gamma(m+n-\alpha) \int_0^{2\pi} \lambda(\vartheta)^{\alpha-m-n} (\cos \vartheta)^m (\sin \vartheta)^n g(\vartheta)d\vartheta.$$

Remark 2. The 1-D projection $Y = v_1X_1 + v_2X_2$ discussed in Remark 1, with $v = \sqrt{v_1^2 + v_2^2}$, gives in the continuous case the cumulant function

$$\begin{aligned} \kappa_Y(y) &= \int_0^{2\pi} \lambda(\vartheta)^\alpha \psi_\alpha\left(\frac{v \cos(\vartheta - \theta_{v_2/v_1})}{\lambda(\vartheta)}y\right)g(\vartheta)d\vartheta \\ &= \int_0^{2\pi} \lambda(\vartheta)^\alpha \psi_\alpha\left(\frac{v \cos(\vartheta - \vartheta')}{\lambda(\vartheta)}y\right)g(\vartheta)d\vartheta, \end{aligned}$$

where $\vartheta' \in [0, 2\pi]$, and $v > 0$ are fixed, with the cumulants

$$\text{Cum}_m(Y) = \Gamma(m-\alpha) \int_0^{2\pi} \lambda(\vartheta)^{\alpha-m} (v \cos(\vartheta - \vartheta'))^m g(\vartheta)d\vartheta.$$

4 Tempering via special functions

The Smoothly Truncated Lévy Distributions were defined in the previous section through an exponential tempering functions which, in turn, determined the measures Q and R via the formula (7). In this section we will review several other examples of tempering via special functions such as gamma, Bessel, alpha-stable density, etc. These models will approximate the stable distribution with different speeds and will have different probabilistic properties expressed by the behavior of their cumulants. All of the models will depend on parameter(s) λ ; as the parameter goes to 0, the model converges to the Lévy stable distribution; some interesting new probability distributions arise in the process as well.

In the following we consider only the one-sided ($p_1 = 0$) measure σ on the unit ball $\mathbb{S}^0 = \{-1, 1\}$. The two sided case is can be handled as in the case of Smoothly Truncated Lévy Distribution described in Subsection 3.3: the two-sided measure is given by the formula

$$\sigma(\{-1\}) = ap_1, \quad \sigma(\{1\}) = ap_2,$$

where $a, p_1, p_2 > 0$, $p_1 + p_2 = 1$, while the tempering does not depend on the direction, i.e. $k(r|\pm 1) = k(r)$. Subsection 3.4 gives also a general method for extending the results of this section to a family of more flexible models.

4.1 Mixed tempering

In this subsection we consider a truncated tempering scheme which permits control of the closeness to the α -stable distribution through two parameters λ_1 and λ_2 . So, let the tempering function $k(r)$ be

$$k(r) = (1 + \lambda_2 r)^{-\beta} \exp(-\lambda_1 r), \quad \lambda_1, \lambda_2 > 0, \quad r > 0.$$

Then, $k(r)$ is the Laplace transform of the function

$$Q(dx) = \frac{(x - \lambda_1)_+^{\beta-1}}{\lambda_2^\beta \Gamma(\beta)} \exp\left(-\frac{[x - \lambda_1]_+}{\lambda_2}\right) dx,$$

so that, in view of (6), the Q -measure

$$Q(B) = \int_0^\infty 1_B(x) \frac{(x - \lambda_1)_+^{\beta-1}}{\lambda_2^\beta \Gamma(\beta)} \exp\left(-\frac{[x - \lambda_1]_+}{\lambda_2}\right) dx,$$

and the R -measure (see, (7)) is given by the formula

$$\begin{aligned} R(B) &= \int_{\mathbb{R}_+} 1_B\left(\frac{\text{sign}(x)}{|x|}\right) |x|^\alpha Q(dx) \\ &= \int_{\lambda_1}^\infty 1_B\left(\frac{1}{x}\right) |x|^\alpha \frac{(x - \lambda_1)^{\beta-1}}{\lambda_2^\beta \Gamma(\beta)} \exp\left(-\frac{x - \lambda_1}{\lambda_2}\right) dx \\ &= \int_0^{1/\lambda_1} 1_B(y) y^{-\beta-\alpha-1} \frac{(1 - \lambda_1 y)^{\beta-1}}{\lambda_2^\beta \Gamma(\beta)} \exp\left(-\frac{1/y - \lambda_1}{\lambda_2}\right) dy. \end{aligned}$$

Since the density ρ of the R -measure

$$\rho(y) = 1_{0 < y < 1/\lambda_1} y^{-\beta-\alpha-1} \frac{(1 - \lambda_1 y)^{\beta-1}}{\lambda_2^\beta \Gamma(\beta)} \exp\left(-\frac{1/y - \lambda_1}{\lambda_2}\right),$$

is zero outside of a finite interval, all the moments, and the moment generating function, exist, see [18]. The cumulants c_m are then calculated from the Lévy measure M , namely

$$\begin{aligned} c_m &= \int_0^\infty r^{m-\alpha-1} (1 + \lambda_2 r)^{-\beta} \exp(-\lambda_1 r) dr \\ &= \lambda_2^{-\beta} \int_0^\infty r^{m-\alpha-1} (1/\lambda_2 + r)^{-\beta} \exp(-\lambda_1 r) dr \\ &= \lambda_2^{-\beta} \Gamma(m - \alpha) \lambda_2^{\beta+\alpha-m} U(m - \alpha, m - \alpha + 1 - \beta, \lambda_1/\lambda_2), \end{aligned}$$

for $m > \alpha$, we have

$$c_m = \lambda_2^{\alpha-m} \Gamma(m - \alpha) U(m - \alpha, m - \alpha + 1 - \beta, \lambda_1/\lambda_2),$$

where U is the Kummer's function of the second kind, also known as the confluent hypergeometric function of the second kind ${}_1F_1$, see [17], 2.3.6.9. Naturally if λ_2 tends to zero then c_m is close to the cumulant of Lévy Flights since $U(m - \alpha, m - \alpha + 1 - \beta, \lambda_1/\lambda_2) = (\lambda_1/\lambda_2)^{\alpha-m} (1 + O(\lambda_2/\lambda_1))$.

4.2 Gamma tempering.

In this subsection the exponential, smoothly truncated tempering is replaced by a slower-decaying Gamma tempering. So, let the tempering function $k(r)$ be

$$k(r) = (1 + \lambda r)^{-\beta}, \quad r > 0.$$

Then $Q(ds|y)$ is the Gamma distribution with the Laplace transform

$$q(r) = \frac{x^{\beta-1}}{\lambda^\beta \Gamma(\beta)} \exp(-x/\lambda).$$

It is straightforward to check that the measures Q and R , (see (6), and (7), respectively) are given by the formulas

$$Q(dx) = 1_{x>0} \frac{x^{\beta-1}}{\lambda^\beta \Gamma(\beta)} \exp(-x/\lambda) dx,$$

and

$$\begin{aligned} R(B) &= \int_{\mathbb{R}_0} 1_B \left(\frac{\text{sign}(x)}{|x|} \right) |x|^\alpha Q(dx) \\ &= \int_0^\infty 1_B \left(\frac{1}{x} \right) \frac{x^{\beta+\alpha-1}}{\lambda^\beta \Gamma(\beta)} \exp(-x/\lambda) dx \\ &= \int_0^\infty 1_B(y) \frac{y^{-\beta-\alpha-1}}{\lambda^\beta \Gamma(\beta)} \exp[-1/(\lambda y)] dy, \end{aligned}$$

The density of the R- measure, with $0 < \beta + \alpha$, is thus

$$\rho(y) = 1_{y>0} \frac{y^{-\beta-\alpha-1}}{\lambda^\beta \Gamma(\beta)} \exp[-1/(\lambda y)],$$

is the reciprocal (or inverse) of the Γ - density. The moments of order $m < \beta + \alpha$ of the R- measure exist, and

$$\mu_{R,m} = \frac{\lambda^{\alpha-m} \Gamma(\alpha + \beta - m)}{\Gamma(\beta)},$$

with the cumulants of the form

$$c_m = \frac{\lambda^{\alpha-m} \Gamma(m - \alpha) \Gamma(\alpha + \beta - m)}{\Gamma(\beta)}.$$

4.3 General Inverse Gaussian tempering.

The Generalized Inverse Gaussian (GIG) distribution has the density

$$q(x) = \frac{(a/b)^{h/2}}{2K_h(\sqrt{ab})} x^{h-1} \exp\left(-\frac{1}{2} [ax + bx^{-1}]\right),$$

where K_h is the modified Bessel function of the second kind with index h . Notice that after replacing the parameters a , and b , by the parameters a/λ , and λb , respectively, the density takes the form

$$q(x) = \frac{(a/b)^{h/2} \lambda^{-h}}{2K_h(\sqrt{ab})} x^{h-1} \exp\left(-\frac{1}{2} [ax/\lambda + \lambda bx^{-1}]\right),$$

where K_h is Modified Bessel Function of the Second Kind, and the parameter λ can now be used for tuning the TS_α distribution close to the α -stable distribution. Let $h < 0$, then the Laplace transform $k(r)$ of $q(x)$, is well known (see Halgreen [7]),

$$k(r) = \exp\left(-\int_{a/2}^{\infty} bg_{-h}(2xb-ab)\ln\left(1+\frac{\lambda r}{x}\right)dx\right),$$

where

$$g_{-h}(x) = 2\left(\pi^2x\left[J_{|h|}^2(\sqrt{x})+N_{|h|}^2(\sqrt{x})\right]\right)^{-1},$$

J_h and N_h are Bessel functions, see [6], Chapter 8. Observe that, for fixed a, b , and h , the limit of $k(r)$, as $\lambda \rightarrow 0$, is equal to 1. The measure Q is concentrated on the half-line $(0, \infty)$, and

$$Q(dx) = q(x)dx,$$

while the corresponding R- measure has the density ρ of the form

$$\rho(x) = \frac{(a/b)^{h/2}\lambda^{-h}}{2K_h(\sqrt{ab})}x^{-\alpha-h-1}\exp\left(-\frac{1}{2}\left[(a/\lambda)x^{-1}+b\lambda x\right]\right),$$

The moments of the R- measure are

$$\begin{aligned}\mu_{R,m} &= \frac{(a/b)^{h/2}\lambda^{-h}}{2K_h(\sqrt{ab})}\int_0^\infty x^{m-\alpha-h-1}\exp\left(-\frac{1}{2}\left[(a/\lambda)x^{-1}+b\lambda x\right]\right)dx \\ &= \left(\frac{a}{b\lambda^2}\right)^{(m-\alpha)/2}\frac{K_{m-\alpha-h}(\sqrt{ab})}{K_h(\sqrt{ab})},\end{aligned}$$

see [6] 3.471.9. In particular, if $h = -1$, and $\alpha = 1/2$, then

$$\mu_{R,m} = \left(\frac{a}{b\lambda^2}\right)^{(m-\alpha)/2}\frac{K_{m+1/2}(\sqrt{ab})}{K_1(\sqrt{ab})},$$

where

$$K_{m+1/2}(y) = \sqrt{\frac{\pi}{2y}}e^{-y}\sum_{k=0}^m\frac{(m+k)!}{k!(m-k)!(2y)^k}.$$

see [6] 3.468. Similar results can be obtained for any integer h , and $\alpha = 1/2$, or $3/2$.

4.4 Fractional exponential.

The function

$$k(r) = \frac{1}{(\lambda r)^\beta + 1},$$

is the Laplace transform of the Fractional Exponential Function $q(x)$. The moments up to order $\alpha + \beta$ exist. In particular for $\beta = 1/2$, we have

$$q(x) = \sqrt{\frac{\pi}{\lambda x}} - \frac{\pi}{\lambda}e^{x/\lambda}\operatorname{erfc}\left(\sqrt{x/\lambda}\right),$$

where erfc is the complementary error function (see [16]). From

$$Q(dx) = q(x) dx,$$

In this case, the R -measure is of the form

$$\begin{aligned} R(B) &= \int_{\mathbb{R}_+} 1_B\left(\frac{1}{x}\right) x^\alpha q(x) dx \\ &= \int_{\mathbb{R}_+} 1_B\left(\frac{1}{x}\right) \left[\sqrt{\frac{\pi}{\lambda x}} - \frac{\pi}{\lambda} e^{x/\lambda} \operatorname{erfc}\left(\sqrt{x/\lambda}\right) \right] x^\alpha dx \\ &= \int_{\mathbb{R}_+} 1_B(x) \left[\sqrt{\frac{\pi x}{\lambda}} - \frac{\pi}{\lambda} e^{1/(x\lambda)} \operatorname{erfc}\left(1/\sqrt{\lambda x}\right) \right] \frac{dx}{x^{2+\alpha}} \end{aligned}$$

with the density

$$\rho(x) = \frac{1}{x^{2+\alpha}} \left[\sqrt{\frac{\pi x}{\lambda}} - \frac{\pi}{\lambda} e^{1/(x\lambda)} \operatorname{erfc}\left(1/\sqrt{\lambda x}\right) \right].$$

Also the verification of the existence of cumulants (moments) of order $m < \alpha + 1/2$ is here straightforward since, for large x , we have the asymptotics

$$\sqrt{\frac{\pi}{x}} - \pi e^x \operatorname{erfc}(\sqrt{x}) \sim \sqrt{\frac{\pi}{4x^3}},$$

see [6] 8.254. For small x ,

$$\sqrt{\frac{\pi}{x}} - \pi e^x \operatorname{erfc}(\sqrt{x}) \sim \sqrt{\frac{\pi}{x}},$$

so that, if $\alpha > 1/2$ then the measure R is not finite. The condition (13) is fulfilled only in some cases so that some assumptions have to be made before formula (14) can be used to calculate the cumulants.

Thus, for $1/2 < \alpha < 1$, the first order cumulant (the expectation)

$$c_1 = \frac{2\pi\lambda^{\alpha-1}}{\sin(2\pi[1-\alpha])},$$

and, for $3/2 < \alpha < 2$, the second order cumulant (the variance)

$$c_2 = \frac{2\pi\lambda^{\alpha-2}}{\sin(2\pi[2-\alpha])}.$$

Of course, one can try to calculate the cumulants directly from the Lévy measure M , but the basic formula

$$c_m = \int_{\mathbb{R}_+} x^m M(dx) = \int_0^\infty \frac{r^{m-\alpha-1}}{\sqrt{\lambda r + 1}} dr,$$

works only for $\alpha < m < \alpha + 1/2$ (since assumption (3) is not satisfied), although we know that c_m exists for all $m < \alpha + 1/2$.

4.5 1/3-stable tempering.

Another example of the tempering function for which explicit calculations are possible is the case when $q(x)$ is the density of the one-sided 1/3-stable distribution, that is

$$q(x) = \frac{\sqrt{\lambda}}{3\pi} x^{-3/2} K_{1/3} \left(\frac{2\sqrt{\lambda}}{3^{3/2}\sqrt{x}} \right), \quad x > 0.$$

where $K_{1/3}(x)$ is the modified Bessel function of the second kind with order 1/3 (see, Subsection 4.3). Its Laplace transform

$$k(r) = \exp \left(-(\lambda r)^{1/3} \right).$$

In this context the R- measure and its density are easily determined as

$$\begin{aligned} R(B) &= \int_{\mathbb{R}_+} 1_B \left(\frac{1}{x} \right) x^\alpha q(x) dx \\ &= \frac{\sqrt{\lambda}}{3\pi} \int_{\mathbb{R}_+} 1_B \left(\frac{1}{x} \right) x^{\alpha-3/2} K_{1/3} \left(\frac{2\sqrt{\lambda}}{3^{3/2}\sqrt{x}} \right) dx \\ &= \frac{\sqrt{\lambda}}{3\pi} \int_{\mathbb{R}_+} 1_B(x) x^{-\alpha-1/2} K_{1/3} \left(\frac{2\sqrt{\lambda}}{3^{3/2}\sqrt{x}} \right) dx, \end{aligned}$$

The R -measure is finite if $\alpha < 1/3$, since its density $\rho(x) \sim x^{(1/3-\alpha)-1}$ around zero. Under the assumption $2m - 2\alpha \pm 1/3 > -1$ (see, [1] 11.4.22), the cumulants are then calculated as moments of the R- measure:

$$\begin{aligned} \mu_{R,m} &= \frac{\sqrt{\lambda}}{3\pi} \int_{\mathbb{R}_+} x^{m-\alpha-1/2} K_{1/3} \left(\frac{2\sqrt{\lambda}}{3^{3/2}\sqrt{x}} \right) dx \\ &= \frac{2\sqrt{\lambda}}{3\pi} \int_{\mathbb{R}_+} x^{2m-2\alpha} K_{1/3} \left(\frac{2\sqrt{\lambda}}{3^{3/2}x} \right) dx \\ &= \frac{2\sqrt{\lambda}}{3\pi} \left(\frac{2\sqrt{\lambda}}{3^{3/2}} \right)^{2\alpha-2m-1} \int_{\mathbb{R}_+} x^{2m-2\alpha} K_{1/3}(x) dx \\ &= \frac{2\lambda^{\alpha-m}}{3\pi} \left(\frac{2}{3^{3/2}} \right)^{2\alpha-2m-1} 2^{2m-2\alpha-1} \Gamma \left(\frac{2m-2\alpha+4/3}{2} \right) \Gamma \left(\frac{2m-2\alpha+2/3}{2} \right) \\ &= \frac{\lambda^{\alpha-m} 3^{1/2-3\alpha+3m}}{2\pi} \Gamma \left(m - \alpha + \frac{2}{3} \right) \Gamma \left(m - \alpha + \frac{1}{3} \right). \end{aligned}$$

The assumption implies that $\alpha - m < 1/3$, so α must be $< 1/3$ for evaluation of the $\mu_{R,m}$, $0 \leq m < \max(0, \alpha - 1/3)$.

For the cumulants of an arbitrary integer order m we have the formula

$$\begin{aligned} c_m &= \Gamma(m - \alpha) \mu_{R,m} \\ &= 3 \frac{\lambda^{\alpha-m} 3^{3(m-\alpha)-1/2}}{2\pi} \Gamma(m - \alpha) \Gamma \left(m - \alpha + \frac{2}{3} \right) \Gamma \left(m - \alpha + \frac{1}{3} \right) \\ &= 3\lambda^{\alpha-m} \Gamma(3[m - \alpha]), \end{aligned}$$

unless $\alpha < 4/3$. If $\alpha > 4/3$, the first cumulant ($m = 1$) can be calculated directly from the Lévy measure:

$$\begin{aligned} c_1 &= \int_{\mathbb{R}_+} xM(dx) = \int_{\mathbb{R}_+} r \exp\left(-(\lambda r)^{1/3}\right) \frac{dr}{r^{\alpha+1}} \\ &= \int_{\mathbb{R}_+} r^{-\alpha} \exp\left(-(\lambda r)^{1/3}\right) dr = 3\lambda^{\alpha-1}\Gamma(3[1-\alpha]), \end{aligned}$$

so that, for all integers m ,

$$c_m = 3\lambda^{\alpha-m}\Gamma(3[m-\alpha]).$$

This is a case when (3) is satisfied but (10) is not.

4.6 Bessel tempering

The function

$$k(r) = \frac{1}{(\lambda r + 1)^{\eta+1}} \exp\left(-\beta \frac{r}{r + 1/\lambda}\right)$$

is the Laplace transform $k(r)$ of the Bessel density

$$q(x) = \lambda^{-1} \exp(-\beta - x/\lambda) \sqrt{\left(\frac{x}{\lambda\beta}\right)^\eta} I_\eta\left(2\sqrt{\beta x/\lambda}\right),$$

where λ, β, η are positive and I_η is a Modified Bessel Function of the First Kind (see, [5], Ch. XIII, Section 3.¹). The corresponding R- measure

$$\begin{aligned} R(B) &= \int_{\mathbb{R}_+} 1_B\left(\frac{1}{x}\right) x^\alpha q(x) dx \\ &= \lambda^{-1-\eta/2} \beta^{-\eta/2} e^{-\beta} \int_{\mathbb{R}_+} 1_B\left(\frac{1}{x}\right) x^{\alpha+\eta/2} \exp(-x/\lambda) I_\eta\left(2\sqrt{\beta x/\lambda}\right) dx \\ &= \lambda^{-1-\eta/2} \beta^{-\eta/2} e^{-\beta} \int_{\mathbb{R}_+} 1_B(x) x^{-\alpha-\eta/2-2} \exp(-1/[\lambda x]) I_\eta\left(2\sqrt{\beta/[\lambda x]}\right) dx. \end{aligned}$$

The limiting behavior of I_η is known ([1] 9.6.7 and 9.7.1) as

$$I_\eta(x) \sim \begin{cases} \left(\frac{1}{2}x\right)^\eta / \Gamma(\eta + 1) & \text{if } x \text{ is small,} \\ \frac{\exp(x)}{\sqrt{2\pi x}} & \text{if } x \text{ is large.} \end{cases}$$

Hence the moments of order $m < \alpha + 3\eta/2 + 1$ exist and the corresponding cumulants can be calculated by formula (14) (see [6] 6.643.2, for the expression in terms of confluent hypergeometric functions).

5 Tempered Stable Ornstein-Uhlenbeck processes

The well known connection between self-decomposable distribution and Ornstein-Uhlenbeck processes is described in, e.g., [20], and [31]. Here, we consider the stationary Ornstein-Uhlenbeck process X_t given by the usual moving average

$$X_t = \int_{-\infty}^t e^{-\gamma(t-s)} dZ_s, \quad (16)$$

¹See also <http://mathworld.wolfram.com/ModifiedBesselFunctionoftheFirstKind.html>

where $\gamma > 0$, and Z_t is usually called the background driving Lévy process (BDLP) for X_t .

In this case the cumulant function κ_{X_t} of X_t is expressed in terms of the cumulant function κ_{Z_1} of Z_t as follows:

$$\kappa_{X_t}(y) = \int_{-\infty}^t \kappa_{Z_1} \left(e^{-\gamma(t-s)} y \right) ds.$$

Throughout this section we shall assume that the second-order moments of Z_s (and X_s) exist and that the processes are centered.

To study the finite dimensional distributions and higher order spectra of X_t assume

$$\mathbb{E} |Z_1|^m < \infty,$$

and denote

$$c_{Z,m} = \text{Cum}_m(Z_1).$$

Then X_t is stationary of order m , i.e., for each t, h_1, \dots, h_{m-1} ,

$$\text{Cum}(X_t, X_{t+h_1}, \dots, X_{t+h_{m-1}}) = \text{Cum}(X_0, X_{h_1}, \dots, X_{t+h_{m-1}}).$$

The Fourier transform of the cumulants gives the m^{th} order spectrum S_m of X_t . In our case, S_m exists and is given by

$$S_m(\omega_1, \omega_2, \dots, \omega_{m-1}) = \frac{c_{Z,m}}{\prod_{j=1}^m (\gamma - i\omega_j)},$$

where $\omega_m = \sum_{j=1}^{m-1} \omega_j$, see [4], and [25]. In particular, if $m = 2$, then the spectrum is

$$S_2(\omega) = \frac{c_{Z,2}}{|\gamma - i\omega|^2}.$$

Notice that

$$S_2(\omega) = \frac{c_{Z,2}}{2\gamma} \int_{-\infty}^{\infty} e^{-i\omega h - \gamma|h|} dh,$$

and conclude that the covariance function of X_t is

$$C_X(h) = \text{Cov}(X_t, X_{t+h}) = e^{-\gamma|h|} \frac{\text{Var}(Z_1)}{2\gamma}.$$

In general, the symmetry of the cumulant implies that the support of the $\text{Cum}(X_0, X_{h_1}, \dots, X_{t+h_{m-1}})$ is the set $\{0 \leq h_1 \leq h_2 \leq \dots \leq h_{m-1}\}$, which directly leads to the result

$$\text{Cum}(X_t, X_{t+h_1}, \dots, X_{t+h_{m-1}}) = e^{-\gamma \sum_{j=1}^{m-1} h_j} \frac{c_{Z,m}}{m\gamma},$$

where $0 \leq h_1 \leq h_2 \leq \dots \leq h_{m-1}$.

The finite dimensional distributions of the stationary Ornstein-Uhlenbeck process X_t are also determined by the BDLP Lévy process Z_t , and γ . Indeed, the joint cumulant function of

$$(X_t, X_{t+h_1}, \dots, X_{t+h_{m-1}})$$

is given by

$$\kappa_{t, t+h_1, \dots, t+h_{m-1}}(y_1, y_2, \dots, y_m) = \kappa_{X_0} \left(y_1 + \sum_{j=1}^{m-1} e^{-\gamma h_j} y_{j+1} \right) + \sum_{j=1}^{m-1} \kappa_{I(\Delta h_j)} \left(\sum_{n=j}^{m-1} y_{n+1} \right),$$

where $0 \leq h_1 \leq h_2 \leq \dots \leq h_{m-1}$, $\Delta h_j = h_{j+1} - h_j$, and

$$I(h) = \int_0^h e^{-\gamma s} dZ_s.$$

The cumulant functions of X_0 and $I(\Delta h_j)$ are obtained from the general formula (12). Hence

$$\begin{aligned} \kappa_{t, t+h_1, \dots, t+h_{m-1}}(y_1, y_2, \dots, y_m) &= \int_0^\infty \kappa_{Z_1} \left(e^{-\gamma s} \left[y_1 + \sum_{j=1}^{m-1} e^{-\gamma h_j} y_{j+1} \right] \right) ds \\ &\quad + \sum_{j=1}^{m-1} \int_0^{\Delta h_j} \kappa_{Z_1} \left(e^{-\gamma s} \sum_{n=j}^{m-1} y_{n+1} \right) ds. \end{aligned}$$

The easy consequence of this formula is that the distribution of X_0 and γ determine the finite dimensional distributions of X_t . The basic example here is the Gaussian Ornstein-Uhlenbeck process: If either X_t , or Z_t , is Gaussian then both Lévy measures M_X and M_Z are zero, $b = 0$ and

$$C_X(h) = e^{-\gamma h} \text{Var}(Z_1) / 2\gamma.$$

All higher order cumulants are zero.

5.1 BDLP for Tempered Stable $TS_\alpha(R, b)$.

From now onwards we concentrate our attention on Ornstein-Uhlenbeck processes for which either X_t , or its BDLP Z_t , is a Gaussian-free, and centered tempered stable processes.

First, consider the case when X_t is a $TS_\alpha(R, 0)$ process with the cumulant function

$$\kappa_X(y) = \int_{\mathbb{R}_0^d} \psi_\alpha(\langle y, x \rangle) R_X(dx),$$

where ψ_α is given by (9). The general formula (19) makes it possible to express the cumulant function κ_Z of the BDLP in terms of measure R_X . A similar result in terms of Lévy densities has been obtained in [3].

Lemma 2 *Let X_t be a $TS_\alpha(R, 0)$ process with the cumulant function*

$$\kappa_X(y) = \int_{\mathbb{R}_0^d} \psi_\alpha(\langle y, x \rangle) R_X(dx),$$

where ψ_α is given by (9). Then, for any $0 < \alpha < 2$, the cumulant function κ_Z of the associated BDLP is given by the formula

$$\kappa_Z(y) = \gamma \int_{\mathbb{R}_0^d} \xi_\alpha(\langle y, x \rangle) R_X(dx), \tag{17}$$

where

$$\xi_\alpha(r) = r \frac{d\psi_\alpha(r)}{dr}.$$

Moreover, if the R -measure of X has a differentiable density ρ_X such that the gradient

$$\nabla_x [x\rho_X(x)] = \left[\frac{\partial}{\partial x_1} x\rho_X(x), \frac{\partial}{\partial x_2} x\rho_X(x), \dots, \frac{\partial}{\partial x_d} x\rho_X(x) \right]$$

is continuous at zero, then the BDLP is a tempered stable process with

$$\kappa_Z(y) = \int_{\mathbb{R}_0^d} \psi_\alpha(\langle y, x \rangle) \rho_Z(x) dx,$$

where the R -density ρ_Z is given by

$$\rho_Z(x) = -\gamma \operatorname{Tr} \nabla_x [\rho_X(x) x] = -\gamma [d\rho_X(x) + [\nabla_x \rho_X(x)] x]. \quad (18)$$

In the one-dimensional case, $d = 1$, we have

$$\rho_Z(x) = -\gamma \left[\rho_X(x) + x \frac{d\rho_X(x)}{dx} \right],$$

and

$$\kappa_Z(y) = \int_{\mathbb{R}_0} \psi_\alpha(yx) \rho_Z(x) dx.$$

Proof. In the polar representation the Lévy measure of X is

$$M(B) = \int_{\mathbb{S}^{d-1}} \sigma(du) \int_0^\infty 1_B(ru) k(r|u) \frac{dr}{r^{\alpha+1}},$$

and the Lévy measure of the BDLP

$$\begin{aligned} N(B) &= -\gamma \int_{\mathbb{S}^{d-1}} \sigma(du) \int_0^\infty 1_B(ru) d \left[\frac{k(r|u)}{r^\alpha} \right] \\ &= -\gamma \int_{\mathbb{S}^{d-1}} \sigma(du) \int_0^\infty 1_B(ru) \left(r \frac{\partial k(r|u)}{\partial r} - \alpha k(r|u) \right) \frac{dr}{r^{\alpha+1}}, \end{aligned} \quad (19)$$

Now, the second term in the last integral

$$\gamma \int_{\mathbb{S}^{d-1}} \sigma(du) \int_0^\infty 1_B(ru) \alpha k(r|u) \frac{dr}{r^{\alpha+1}},$$

provides the representation (8). The representation of the first term containing $\partial k(r|u)/\partial r$ needs some extra work. We have

$$\frac{\partial k(r|u)}{\partial r} = - \int_0^\infty s e^{-rs} Q(ds|u),$$

so that

$$\begin{aligned} \int_{\mathbb{S}^{d-1}} \sigma(du) \int_0^\infty 1_B(ru) r \frac{\partial k(r|u)}{\partial r} \frac{dr}{r^{\alpha+1}} &= - \int_{\mathbb{S}^{d-1}} \sigma(du) \int_0^\infty 1_B(ru) r \int_0^\infty s e^{-rs} Q(ds|u) \frac{dr}{r^{\alpha+1}} \\ &= - \int_{\mathbb{S}^{d-1}} \sigma(du) \int_0^\infty \left(\int_0^\infty 1_B\left(\frac{t}{s}u\right) e^{-t} t^{-\alpha} dt \right) s^\alpha Q(ds|u) \\ &= - \int_0^\infty \int_{\mathbb{R}_0} 1_B\left(t \frac{x}{\|x\|^2}\right) \|x\|^\alpha Q(dx) e^{-t} t^{-\alpha} dt \\ &= - \int_{\mathbb{R}_0} \int_0^\infty 1_B(tx) e^{-t} t^{-\alpha} dt R(dx). \end{aligned}$$

Proceedings in the footsteps of Rosiński's result [18], we notice that , for $0 < \alpha < 1$,

$$\alpha\psi_\alpha(r) + \psi_{\alpha-1}(r) = ir\Gamma(1-\alpha)(1-ir)^{\alpha-1},$$

and, for $1 < \alpha < 2$,

$$\alpha\psi_\alpha(r) + \psi_{\alpha-1}(r) = ir\Gamma(1-\alpha)\left[(1-ir)^{\alpha-1} - 1\right].$$

Hence, in both cases we obtain

$$\alpha\psi_\alpha(r) + \psi_{\alpha-1}(r) = r\frac{d\psi_\alpha(r)}{dr}.$$

Note that the case $\alpha - 1 < 0$ does not create any difficulties here. Denote

$$\xi_\alpha(s) = s\frac{d\psi_\alpha(s)}{ds},$$

and observe that $\xi_\alpha(s)$ has the limit zero at zero. Utilizing (8) we obtain the cumulant function

$$\kappa_Z(y) = \gamma \int_{\mathbb{R}_0^d} \alpha\psi_\alpha(\langle y, x \rangle) + \psi_{\alpha-1}(\langle y, x \rangle) R(dx) = \gamma \int_{\mathbb{R}_0^d} \xi_\alpha(\langle y, x \rangle) R(dx).$$

In the one-dimensional case, $d = 1$,

$$\kappa_Z(y) = \gamma \int_{\mathbb{R}_0} \xi_\alpha(yx) \rho_X(x) dx = \gamma \int_{\mathbb{R}_0} \frac{d\psi_\alpha(r)}{dr} \Big|_{r=yx} yx \rho_X(s) ds,$$

and splitting the domain of integration into $(-\infty, 0)$ and $(0, \infty)$, and integrating by parts,

$$\int_0^\infty \frac{d\psi_\alpha(yx)}{dx} x \rho_X(x) dx = - \int_0^\infty \psi_\alpha(yx) \frac{d}{dx} [x \rho_X(x)] dx.$$

we obtain

$$\frac{d}{dx} x \rho_X(x) = \rho_X(x) + x \frac{d\rho_X(x)}{dx}.$$

Therefore, putting

$$\rho_Z(x) = -\gamma \left[\rho_X(x) + x \frac{d\rho_X(x)}{dx} \right],$$

we see that the density ρ_Z of the R- measure of Z satisfies the equation

$$\kappa_Z(y) = \int_{\mathbb{R}_0} \psi_\alpha(yx) \rho_Z(x) dx.$$

Now , we prove the case $d = 2$; in the general case $d > 2$ the argument is almost the same, the only difference being that the factor r^{d-1} has to be replaced by the Jacobian. So, we have

$$\kappa_Z(y) = -\gamma \int_{\mathbb{R}_0^d} \xi_\alpha(\langle y, x \rangle) R(dx) = -\gamma \int_{\mathbb{R}_0^d} \xi_\alpha(\langle y, x \rangle) \rho_X(x) dx.$$

Fix a $y \neq 0$, and perform a rotation so that the first axis points in the direction of y , and change the integral to the polar coordinate system:

$$\begin{aligned} \int_{\mathbb{R}_0^2} \xi_\alpha(\langle y, x \rangle) \rho_X(x) dx &= \int_0^\infty \int_0^{2\pi} \xi_\alpha(\|y\| r \cos \omega) \rho_X(r \cos \omega, r \sin \omega) r dr d\omega \\ &= \int_0^\infty \int_0^{2\pi} \|y\| r \cos \omega \frac{d\psi_\alpha(s)}{ds} \Big|_{s=\|y\| r \cos \omega} \rho_X(r \cos \omega, r \sin \omega) r dr d\omega \\ &= \int_0^\infty \int_0^{2\pi} \frac{d\psi_\alpha(\|y\| r \cos \omega)}{dr} \rho_X(r \cos \omega, r \sin \omega) r^2 dr d\omega. \end{aligned}$$

Integrating by parts we obtain

$$\begin{aligned} & - \int_0^\infty \int_0^{2\pi} \frac{d\psi_\alpha(\|y\| r \cos \omega)}{dr} \rho_X(r \cos \omega, r \sin \omega) r^2 dr d\omega = \int_0^\infty \int_0^{2\pi} \psi_\alpha(\|y\| r \cos \omega) \\ & \times \left[2\rho_X(r \cos \omega, r \sin \omega) + r \cos \omega \frac{\partial}{\partial x_1} \rho_X(r \cos \omega, r \sin \omega) + r \sin \omega \frac{\partial}{\partial x_2} \rho_X(r \cos \omega, r \sin \omega) \right] r dr d\omega \\ & = \int_{\mathbb{R}_0^2} \psi_\alpha(\langle y, x \rangle) \{2\rho_X(x) + [\nabla_x \rho_X(x)] x\} dx, \end{aligned}$$

and (18) follows.

If the R- density of X is differentiable then the calculation of κ_Z is straightforward; otherwise we have to use the formula (17).

Example 1 Let X be $STLF_{1/2}(a, 0, \lambda)$ with the cumulant function

$$\kappa_X(u) = -2a\sqrt{2\pi\lambda} \left[\sqrt{1 - iu/\lambda} - 1 \right].$$

see Section 3.3. Then the cumulant function of the BDLP Z is

$$\kappa_Z(u) = ia\gamma u (1 - iu/\lambda)^{-1/2}.$$

5.2 Stationary Multivariate Ornstein–Uhlenbeck with Tempered Stable BDLP

Consider now a stationary Ornstein–Uhlenbeck process X_t with parameter γ and BDLP Z_t , where $Z_1 \in TS_\alpha(R_Z, 0)$. The cumulant function κ_{Z_1} of the BDLP is given in terms of the R -measure:

$$\kappa_{Z_1}(y) = \int_{\mathbb{R}_0^d} \psi_\alpha(\langle y, x \rangle) R_Z(dx),$$

see (9) for the definition of the function ψ_α . For each t , the cumulant function of X_t can be written in the form

$$\begin{aligned} \kappa_{X_t}(y) &= \int_{-\infty}^t \kappa_{Z_1}(e^{-\gamma(t-s)}y) ds \\ &= \int_{-\infty}^t \int_{\mathbb{R}_0^d} \psi_\alpha(\langle e^{-\gamma(t-s)}y, x \rangle) R_Z(dx) ds \\ &= \int_{\mathbb{R}_0^d} \int_0^\infty \psi_\alpha(\langle e^{-\gamma s}y, x \rangle) ds R_Z(dx). \end{aligned}$$

Now, formula (11), and Property 3 imply

$$\kappa_{X_t}(y) = \int_{\mathbb{R}_0^d} \psi_\alpha(\langle y, x \rangle) R_X(dx),$$

with the R -measure

$$R_X(B) = \int_0^\infty R_Z(e^{\gamma s} B) ds = \frac{1}{\gamma} \int_1^\infty R_Z(sB) \frac{ds}{s}.$$

Hence X_t is Tempered Stable. If $d = 1$ it follows that the R -density $\rho_X(x)$ exists and

$$\rho_X(x) = \begin{cases} \frac{1}{\gamma x} R_Z([x, \infty)), & x > 0, \\ \frac{1}{\gamma |x|} R_Z((-\infty, x]), & x < 0. \end{cases}$$

Hence $\rho_X(x)$ fulfils the equation

$$\rho_Z(x) = -\gamma \frac{d}{dx} [x \rho_X(x)], \quad (20)$$

as long as ρ_X is differentiable and $\frac{d}{dx} [x \rho_X(x)]$ is continuous at zero.

Example 2 Let Z_1 be an STLTF with the R -measure

$$R_Z = \lambda^\alpha a (p_1 \delta_{-1/\lambda} + p_2 \delta_{1/\lambda}).$$

Here (20) does not apply, and we use (11) instead. The R -measure of the stationary Ornstein–Uhlenbeck process X_t with parameter γ is

$$\begin{aligned} R_X(B) &= \frac{1}{\gamma} \int_{\mathbb{R}_0} \int_0^1 1_B(sx) \frac{ds}{s} R_Z(dx) \\ &= \frac{\lambda^\alpha a}{\gamma} \int_0^1 p_1 1_B\left(-\frac{s}{\lambda}\right) + p_2 1_B\left(\frac{s}{\lambda}\right) \frac{ds}{s} \\ &= \frac{\lambda^{\alpha+1} a}{\gamma} \int_0^\lambda p_1 1_B(-s) + p_2 1_B(s) \frac{ds}{s}. \end{aligned}$$

Hence the R -density of X_t is

$$\rho_X(x) = \frac{\lambda^{\alpha+1} a}{\gamma x} (p_2 1_{0 < x \leq \lambda} - p_1 1_{-\lambda \leq x < 0}).$$

Below, we generalize the above result for a stationary multivariate Ornstein–Uhlenbeck process X_t . Let X_t be given by the stochastic differential equation system

$$dX_t = -QX_t dt + dZ_t, \quad (21)$$

where Q is a real, $d \times d$ -matrix such that the real parts of all eigenvalues are positive ($Q \in \mathcal{M}_+(\mathbb{R}^d)$). Then it is well known, see [21], [23], [22], that there exists a stationary solution X_t of the equation (21), which is given by

$$X_t = \int_{-\infty}^t e^{-Q(t-s)} dZ_s,$$

with the cumulant function

$$\kappa_X(u) = \int_0^\infty \kappa_{Z_1}(e^{-sQ^\top} u) ds. \quad (22)$$

i.e. the cumulant function of X_t is determined by that of Z_1 . This result should be compared with those of [2] as well.

Theorem 1 Suppose $Z_1 \in TS_\alpha(R_Z, 0)$, then the multivariate stationary Ornstein–Uhlenbeck process X_t , with matrix $Q \in \mathcal{M}_+(\mathbb{R}^d)$, and BDLP Z_t , is Tempered Stable. The R –measure of X_t is

$$R_X(B) = \int_0^\infty R_Z(e^{Qs}B) ds.$$

The cumulant function

$$\kappa_X(y) = \int_{\mathbb{R}_0^d} \eta_\alpha(y, x, Q) R_Z(dx),$$

where $\eta_\alpha(y, x, Q)$ is given by the equation

$$\eta_\alpha(y, x, Q) = \int_0^\infty \psi_\alpha(\langle e^{-sQ^\top} y, x \rangle) ds.$$

For each $x, y \in \mathbb{R}_0^d$, $\eta_\alpha(y, x, Q)$ fulfils the equation

$$\psi_\alpha(\langle y, x \rangle) = [\nabla_y \eta_\alpha(y, x, Q)] Q^\top y.$$

Moreover, if the R –measure of Z has density ρ_Z , and the differential equation

$$\begin{aligned} \rho_Z(y) &= -\text{Tr} \nabla_y [\rho_X(y) Q^\top y] \\ &= -\rho_X(y) \text{Tr} Q - [\nabla_y \rho_X(y)] Q^\top y \end{aligned}$$

has a solution ρ_X such that $\nabla_x [\rho_X(x) x]$ continuous at zero, then X_t has R –density ρ_X . The cumulants of X_t are given in terms of the cumulants of Z_1 by the formula

$$\underline{\text{Cum}}_m(\underline{X}_t) = \sum_{k=1}^m \left(I^{\otimes(k-1)} \otimes (Q^\top)^{-1} \otimes I^{\otimes(m-k)} \right) \underline{\text{Cum}}_m(\underline{Z}_1),$$

as long as $\underline{\text{Cum}}_m(\underline{Z}_1)$ exists.

Proof. Start with the equation (22)

$$\begin{aligned} \kappa_X(y) &= \int_0^\infty \kappa_{Z_1}(e^{-sQ^\top} y) ds \\ &= \int_0^\infty \int_{\mathbb{R}_0^d} \psi_\alpha(\langle e^{-sQ^\top} y, x \rangle) R_Z(dx) ds \\ &= \int_{\mathbb{R}_0^d} \int_0^\infty \psi_\alpha(x^\top e^{-sQ^\top} y) ds R_Z(dx). \end{aligned}$$

Denote

$$\eta_\alpha(y, x) = \int_0^\infty \psi_\alpha(x^\top e^{-sQ^\top} y) ds,$$

and apply the gradient operator ∇_y

$$\nabla_y \eta_\alpha(y, x) = \int_0^\infty \frac{d\psi_\alpha(r)}{dr} \Big|_{r=x^\top e^{-sQ^\top} y} x^\top e^{-sQ^\top} ds.$$

Now, let us differentiate $\psi_\alpha(x^\top e^{-Qs}y)$ with respect to s ,

$$\frac{d\psi_\alpha(x^\top e^{-sQ^\top} y)}{ds} = - \frac{d\psi_\alpha(r)}{dr} \Big|_{r=x^\top e^{-sQ^\top} y} x^\top e^{-sQ^\top} Q^\top y,$$

and integrate,

$$\begin{aligned}\psi_\alpha(\langle y, x \rangle) &= \int_0^\infty \frac{d\psi_\alpha(r)}{dr} \Big|_{r=x^\top e^{-sQ^\top} y} x^\top e^{-sQ^\top} ds Q^\top y \\ &= [\nabla_y \eta_\alpha(y, x)] Q^\top y,\end{aligned}$$

assuming $Q^\top u \neq 0$. Let us consider

$$\begin{aligned}\int_{\mathbb{R}_0^d} \eta_\alpha(y, x) \text{Tr} \nabla_y [\rho_X(y) Q^\top y] dy &= -\text{Tr} \int_{\mathbb{R}_0^d} [\nabla_y \eta_\alpha(y, x)] Q^\top y \rho_X(y) dy \\ &= -\int_{\mathbb{R}_0^d} \psi_\alpha(\langle y, x \rangle) \rho_X(y) dy.\end{aligned}$$

Now we conclude that

$$\begin{aligned}\rho_Z(y) &= -\text{Tr} \nabla_y [\rho_X(y) Q^\top y] \\ &= -\rho_X(y) \text{Tr} Q - [\nabla_y \rho_X(y)] Q^\top y.\end{aligned}$$

Note that the multivariate stationary Ornstein-Uhlenbeck process X_t is Q -decomposable, see Masuda (2004) [14] for details. The cumulants for X_t are

$$\begin{aligned}D_{\underline{u}}^{\otimes k} \kappa_X(y) &= \int_{\mathbb{R}_0^d} D_{\underline{u}}^{\otimes k} \eta_\alpha(y, x) R(dx), \\ D_{\underline{u}}^{\otimes} \kappa_X(y) &= \int_{\mathbb{R}_0^d} \int_0^\infty D_{\underline{u}}^{\otimes} \psi_\alpha(x^\top e^{-sQ^\top} y) ds R(dx) \\ &= \int_{\mathbb{R}_0^d} \int_0^\infty \frac{d\psi_\alpha(r)}{dr} \Big|_{r=x^\top e^{-Qs} y} e^{-sQ^\top} ds x R(dx).\end{aligned}$$

Putting $y = 0$ we obtain

$$\begin{aligned}\mathbf{E}X_t &= \int_0^\infty e^{-sQ^\top} ds \mathbf{E}Z_1 \\ &= (Q^\top)^{-1} \mathbf{E}Z_1.\end{aligned}$$

Now

$$\begin{aligned}D_{\underline{u}}^{\otimes 2} \kappa_X(y) &= \int_{\mathbb{R}_0^d} \int_0^\infty D_{\underline{u}}^{\otimes 2} \psi_\alpha(x^\top e^{-sQ^\top} y) ds R(dx) \\ &= \int_{\mathbb{R}_0^d} \int_0^\infty D_{\underline{u}}^{\otimes} \left[e^{-sQ^\top} x \frac{d\psi_\alpha(r)}{dr} \Big|_{r=x^\top e^{-Qs} y} \right] ds R(dx) \\ &= \int_{\mathbb{R}_0^d} \int_0^\infty [e^{-sQ^\top} x \otimes I] \frac{d^2 \psi_\alpha(r)}{dr^2} \Big|_{r=x^\top e^{-Qs} y} e^{-sQ^\top} x ds R(dx) \\ &= \int_{\mathbb{R}_0^d} \int_0^\infty [e^{-sQ^\top} \otimes e^{-sQ^\top}] \frac{d^2 \psi_\alpha(r)}{dr^2} \Big|_{r=x^\top e^{-Qs} y} x^{\otimes 2} ds R(dx).\end{aligned}$$

Again, putting $u = 0$, we obtain

$$\underline{\text{Cum}}_2(\underline{X}_t) = \left[(Q^\top)^{-1} \otimes I + I \otimes (Q^\top)^{-1} \right] \underline{\text{Cum}}_2(\underline{Z}_1),$$

since

$$\int_0^\infty [e^{-sQ^\top} \otimes e^{-sQ^\top}] ds = (Q^\top)^{-1} \otimes I + I \otimes (Q^\top)^{-1}.$$

The general case $\underline{\text{Cum}}_m(\underline{X}_t)$ is then obtained by a standard induction. ■

6 Appendix

6.1 Multiple cumulants

Introduce the notation $\frac{\partial}{\partial \underline{u}^\top} = \nabla_{\underline{u}}$. The operator $D_{\underline{u}}^{\otimes}$ is defined as

$$D_{\underline{u}}^{\otimes} \underline{\phi} = \text{Vec} \left(\frac{\partial \underline{\phi}}{\partial \underline{u}^\top} \right)^\top = \text{Vec} \begin{bmatrix} \frac{\partial \phi_1}{\partial u_1} & \frac{\partial \phi_1}{\partial u_2} & \dots & \frac{\partial \phi_1}{\partial u_d} \\ \frac{\partial \phi_2}{\partial u_1} & \ddots & & \vdots \\ \vdots & & \ddots & \vdots \\ \frac{\partial \phi_m}{\partial u_1} & \dots & \dots & \frac{\partial \phi_m}{\partial u_d} \end{bmatrix}^\top,$$

which is a column vector of order md . We can also write $D_{\underline{u}}^{\otimes}$ in terms of the Kronecker product.

$$\begin{aligned} D_{\underline{u}}^{\otimes} \underline{\phi} &= \text{Vec} \left(\underline{\phi} \frac{\partial}{\partial \underline{u}^\top} \right)^\top = \text{Vec} \left(\frac{\partial}{\partial \underline{u}} \underline{\phi}^\top \right) \\ &= [\phi_1(\underline{u}), \phi_2(\underline{u}), \dots, \phi_m(\underline{u})]^\top \otimes \left[\frac{\partial}{\partial u_1}, \frac{\partial}{\partial u_2}, \dots, \frac{\partial}{\partial u_d} \right]^\top. \end{aligned}$$

If we repeat the differentiation $D_{\underline{u}}^{\otimes}$ twice, we obtain

$$\begin{aligned} D_{\underline{u}}^{\otimes 2} \underline{\phi} &= D_{\underline{u}}^{\otimes} (D_{\underline{u}}^{\otimes} \underline{\phi}) = \text{Vec} \left[\left(\underline{\phi} \otimes \frac{\partial}{\partial \underline{u}} \right) \frac{\partial}{\partial \underline{u}^\top} \right]^\top \\ &= \underline{\phi} \otimes \left(\frac{\partial}{\partial \underline{u}} \right)^{\otimes 2} = \underline{\phi} \otimes \frac{\partial}{\partial \underline{u}^{\otimes 2}}, \end{aligned}$$

and in general (assuming the existence of derivatives of order k), the k^{th} derivative is given by

$$D_{\underline{u}}^{\otimes k} \underline{\phi} = D_{\underline{u}}^{\otimes} \left(D_{\underline{u}}^{\otimes k-1} \underline{\phi} \right) = [\phi_1(\underline{u}), \phi_2(\underline{u}), \dots, \phi_m(\underline{u})]^\top \otimes \left[\frac{\partial}{\partial u_1}, \frac{\partial}{\partial u_2}, \dots, \frac{\partial}{\partial u_d} \right]^\top{}^{\otimes k}.$$

Let $\varphi(y)$ be the characteristic function of the random variable \underline{X} . Then the multiple moment of order k is defined by

$$\mu_{\otimes j} = (-i)^k D_y^{\otimes k} \varphi(y) \Big|_{y=0},$$

and the multiple k^{th} -order cumulant is

$$c_k = \underline{\text{Cum}}_k(\underline{X}) = (-i)^k D_y^{\otimes k} \log \varphi(y) \Big|_{y=0},$$

see [26] for details. Note that

$$\underline{\text{Cum}}_2(\underline{X}_1, \underline{X}_2) = \mathbb{E}[(\underline{X}_1 - \mathbb{E}\underline{X}_1) \otimes (\underline{X}_2 - \mathbb{E}\underline{X}_2)] = \text{Vec Cov}(\underline{X}_2, \underline{X}_1). \quad (23)$$

References

- [1] Milton Abramowitz and Irene A. Stegun, editors. *Handbook of mathematical functions with formulas, graphs, and mathematical tables*. Dover Publications Inc., New York, 1992. Reprint of the 1972 edition.

- [2] O. E. Barndorff-Nielsen, J. Pedersen, and K. Sato. Multivariate subordination, self-decomposability and stability. *Adv. in Appl. Probab.*, 33(1):160–187, 2001.
- [3] O. E. Barndorff-Nielsen and N. Shephard. Non-Gaussian Ornstein-Uhlenbeck-based models and some of their uses in financial economics. *J. R. Stat. Soc. Ser. B Stat. Methodol.*, 63(2):167–241, 2001.
- [4] D. R. Brillinger. *Time Series; Data Analysis and Theory*. Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, 2001. Reprint of the 1981 edition.
- [5] W. Feller. *An Introduction of Probability Theory and its Application*, volume II. John Wiley, New York, London, 1966.
- [6] I. S. Gradshteyn and I. M. Ryzhik. *Table of integrals, series, and products*. Academic Press Inc., San Diego, CA, sixth edition, 2000. Translated from the Russian, Translation edited and with a preface by Alan Jeffrey and Daniel Zwillinger.
- [7] C. Halgreen. Self-decomposability of the generalized inverse Gaussian and hyperbolic distributions. *Z. Wahrsch. Verw. Gebiete*, 47(1):13–17, 1979.
- [8] P. Hougaard. Survival models for heterogeneous populations derived from stable distributions. *Biometrika*, 73(2):387–396, 1986.
- [9] Z. J. Jurek and W. Vervaat. An integral representation for self-decomposable Banach space valued random variables. *Z. Wahrsch. Verw. Gebiete*, 62(2):247–262, 1983.
- [10] I. Koponen. Analytic approach to the problem of convergence of truncated Lévy flights towards the gaussian stochastic process. *Phys. Rev. E.*, 52:1197–1199, 1995.
- [11] E. Lukacs. A characterization of stable processes. *J. Appl. Probability*, 249:409–418, 1969.
- [12] B.B. Mandelbrot, Variables et processus stochastiques de Pareto-Lévy et la répartition des revenus, I et II. *Comptes Rendus (Paris)*, 6:613–615, 1959.
- [13] R. N. Mantegna and H. E. Stanley. Stochastic processes with ultraslow convergence to a gaussian: The truncated Lévy flight. *Phys. Rev. Lett.*, 73:2946–2949, 1994.
- [14] E.W. Montroll and H. Scher. On multidimensional Ornstein-Uhlenbeck processes driven by a general Lévy process. *Bernoulli*, 10(1):97–120, 2004.
- [15] H. Masuda. Random walks on lattices. IV. Continuous-time walks and influence of absorbing boundaries. *Journal of Statistics Physics*, 9(2):101-135, 1973.
- [16] A. Piryatinska, A.I. Saichev, and W.A. Woyczynski. Models of anomalous diffusion: The subdiffusive case. *Physica A: Statistical Physics*, 349:375–424, 2005.
- [17] A. P. Prudnikov, Y. A. Brychkov I. Marichev. Integrals and series. Vol. 1. Gordon & Breach Science Publishers, New York 1986. Elementary functions, Translated from the Russian and with a preface by N. M. Queen.
- [18] J. Rosinski. Tempering stable processes. Technical report, Department of Mathematics of the Univ. of Tennessee in Knoxville, Tennessee, 2004.

- [19] J. Rosiński. Tempered stable processes. In *2nd MaPhySto Levy Conference*, page 215. MaPhySto, Aarhus, January 2002.
- [20] K. Sato. *Lévy processes and infinitely divisible distributions*, volume 68 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 1999. Translated from the 1990 Japanese original, Revised by the author.
- [21] K. Sato and M. Yamazato. Stationary processes of Ornstein-Uhlenbeck type. In *Probability theory and mathematical statistics (Tbilisi, 1982)*, volume 1021 of *Lecture Notes in Math.*, pages 541–551. Springer, Berlin, 1983.
- [22] K. Sato and M. Yamazato. Operator-self-decomposable distributions as limit distributions of processes of Ornstein-Uhlenbeck type. *Stochastic Process. Appl.*, 17(1):73–100, 1984.
- [23] K. Sato and M. Yamazato. Completely operator-self-decomposable distributions and operator-stable distributions. *Nagoya Math. J.*, 97:71–94, 1985.
- [24] V. Seshadri. *The inverse Gaussian distribution*, volume 137 of *Lecture Notes in Statistics*. Springer-Verlag, New York, 1999. Statistical theory and applications.
- [25] Gy. Terdik. *Bilinear Stochastic Models and Related Problems of Nonlinear Time Series Analysis; A Frequency Domain Approach*, volume 142 of *Lecture Notes in Statistics*. Springer Verlag, New York, 1999.
- [26] Gy. Terdik. Higher order statistics and multivariate vector hermite polynomials for nonlinear analysis of multidimensional time series. *Teor. Ver. Matem. Stat. (Teor. Imovirnost. ta Matem. Statyst.)*, 66:147–168, 2002.
- [27] Gy. Terdik, W. A. Woyczynski, and A. Piryatinska. Fractional- and integer-order moments, and multiscaling for smoothly truncated lévy flights. *Physics Letters A*, 348:94–109, 2006.
- [28] K. Urbanik, Self-decomposable probability distributions on R^m , *Zastos. Mat.* **10** (1969), 91–97.
- [29] K. Urbanik, *Lévy's probability measures on Euclidean spaces*, *Studia Math.* **44** (1972), 119–148, Collection of articles honoring the completion by Antoni Zygmund of 50 years of scientific activity, II.
- [30] K. Urbanik, *Operator-decomposable distributions on Euclidean spaces*, Transactions of the Sixth Prague Conference on Information Theory, Statistical Decision Functions, Random Processes (Tech. Univ. Prague, Prague, 1971; dedicated to the memory of Antonín Špaček), Academia, Prague, 1973, pp. 859–872.
- [31] S. J. Wolfe. On a continuous analogue of the stochastic difference equation $X_n = \rho X_{n-1} + B_n$. *Stochastic Process. Appl.*, 12(3):301–312, 1982.